


CA09/00850

PCT/CA/ 00/00850
14 NOV 2000 (14.11.00)

REC'D 28 NOV 2000	
WIPO	PCT

0/031449

PA 319496



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

October 24, 2000

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/144,812

FILING DATE: July 21, 1999

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)



By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS

P. R. Grant

P. R. GRANT

Certifying Officer

PROVISIONAL APPLICATION COVER SHEET

To the Commissioner of Patents and Trademarks
Washington, DC 20231

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53(b) (2).

Docket No.	13178PRO	Type a plus sign (+) inside this box -	+
------------	----------	--	---

INVENTOR (s) / APPLICANT (s)

Inventor No. 1 of 3	Address:	Citizenship
Last Name: Yudin First Name: Andrei M.I. Konstantin	30 Humewood Drive Toronto, Ontario M6C 2WR Canada	Russian Federation
Inventor No. 2 of 3	Address	Citizenship
Last Name: Martyn First Name: Leo James M.I. Patrick	537 Fairview Rd. W. Mississauga, Ontario L5B 3X1 Canada	Canada
Inventor No. 3 of 3	Address	Citizenship
Last Name: Pandiaraju First Name: Subramanian M.I.	80 St. George St. Toronto, Ontario M5S 3H6 Canada	India

TITLE OF THE INVENTION (280 characters max)

POLYFLUORINATED CATALYSTS

CORRESPONDENCE ADDRESS

Ralph A. Dowell PTO CUSTOMER NO. 000293
DOWELL & DOWELL, P. C.
Suite 309
1215 Jefferson Davis Highway
Arlington, Virginia 22202-3697
Telephone (703) 415-2555
Fax No. (703) 415-2559

STATE	Virginia	ZIP CODE	22202	COUNTRY	United States
-------	----------	----------	-------	---------	---------------

ENCLOSED APPLICATION PARTS (check all that apply)

<input checked="" type="checkbox"/> Specification	Number of Pages	12	<input type="checkbox"/> Small Entity Statement
<input checked="" type="checkbox"/> Drawing(s)	Number of Sheets	1	<input type="checkbox"/> Other (specify)

3654 U.S. PTO

07/21/99

1c503 U.S. PTO
60/144812

07/21/99

60144812-072199





FIG. 1

metal Catalyst using Fluorinated BINOL
Based Catalyst



METHOD OF PAYMENT (check one)

<input checked="" type="checkbox"/> A check or money order enclosed to cover the Provisional filing fees <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees and credit any overages to Deposit Account Number: <u>04-1577</u>	Provisional filing fee amount (s)	150.00
---	-----------------------------------	--------

Respectfully submitted,

Signature: *Ralph A. Dowell*

Date: July 21, 1999

Typed or Printed Name: Ralph A. Dowell Registration No.: 26,868

001220-21844109

07/21/1999

16:33

HILL & SCHUMACHER → DOWELL

NO.815

P03

UNITED STATES

Title: POLYFLUORINATED CATALYSTS

Inventors: Andrei Konstantin Yudin, Leo James Patrick Martyn and
Subramanian Pandiaraju

60144212-072199

07/21 '99 16:32

FIELD OF THE INVENTION

The present invention relates to the design and method of synthesis of asymmetric catalysts and more particularly the present invention provides a new generation of polyfluorinated catalysts.

BACKGROUND OF THE INVENTION

Modern asymmetric synthesis often calls for catalytic transformations. Many important discoveries in this area are due to serendipity, however, understanding the balance of steric and electronic factors is required in order to fine-tune a catalyst toward optimal rate/selectivity in a particular reaction. The analysis of steric environments around metal centers has traditionally dominated attempts to explain and predict the outcome of metal-based enantioselective processes. In comparison, the importance of electronic effects in asymmetric induction was appreciated only in recent years. Several catalytic systems known to date employ electronically diverse substituents on ligands in order to modulate reactivity of the metal center.¹ For example, in the catalytic asymmetric epoxidation of unfunctionalized olefins, electronic properties of substituents on chiral *salen* ligands determine the nature of transition state.^{1a} The later transition state leads to higher enantioselectivities and electronic attenuation of electrophilic Mn=O centers affords higher levels of enantiomeric excess. Enhancement of enantioselectivity through incorporation of fluorine atoms on chiral phosphine ligands in the asymmetric hydrocyanation of olefins was documented.^{1b} The concept of induced electronic asymmetry allows one to increase the enantioselectivity of rhodium-catalyzed hydroboration of olefins.^{1c} Over the years, several modifications of the BINOL skeleton, aimed at changing its properties, have been reported.³ For example, partially

hydrogenated BINOL was used as a catalyst precursor in enantioselective alkylation of aldehydes,^{3a} conjugate addition of diethylzinc to cyclic enones,^{3b} and ring opening of epoxides.^{3c} Incorporation of bromines into the 6 and 6' positions of BINOL, rather remote from the catalytic site, was shown to increase the enantioselectivity of the corresponding titanium catalysts in glyoxal-ene reactions.^{3d}


SUMMARY OF THE INVENTION

It is an object of the present invention to provide a method for designing asymmetric catalysts derived from perfluoroarene ligands.

The present invention provides a new generation of asymmetric polyfluorinated catalysts. Fluorination of aromatic groups on a series of ligands drastically changes their properties including configurational stability and catalytic activity. The fundamental issue is the nature of steric and electronic effects of aromatic fluorination on binaphthol-based catalysts. The basic premise is that alteration of stabilizing stacking and edge-face interactions significantly affects approach of certain substrates to catalytic reaction centers. Due to the high electronegativity of fluorine, electron density in fluoronaphthyl rings is located at the periphery, rather than in the center. The present invention will be illustrated by examples such as preparation of enantiomerically pure fluorobinaphthol ligands and their application in catalytic asymmetric processes.


In one aspect of the invention there is provided method of producing asymmetric polyfluorinated catalysts, comprising: fluorinating aromatic groups on binaphthol-based catalysts.

To prove these points, we prepared racemic form of 2 according to Scheme 1. Tetrafluorobenzene, formed by treating commercially available chloropentafluorobenzene with *n*-butyllithium at -78°C, was reacted with 3-methoxythiophene, obtained from 3-bromothiophene using a literature procedure.⁵ Upon the *in situ* extrusion of sulfur, 2-methoxy-5,6,7,8-tetrafluoronaphthalene 3 was obtained in x% yield. 5,6,7,8-Tetrafluoro-2-naphthol, prepared from 3 by demethylation with BBr₃, did not undergo the FeCl₃-catalyzed oxidative coupling, commonly used for the preparation of 1 from 2-naphthol.⁶ Instead, substitution of hydrogen for chlorine at the 1 position of the aromatic ring took place. Higher oxidation potential of 5,6,7,8-tetrafluoro-2-naphthol (xV vs Ag/AgCl compared to xV vs Ag/AgCl for 1) is a likely reason for the lack of reactivity in the oxidative coupling. We therefore resorted to the reductive route through intermediacy of the 1-brominated derivative 4, prepared in 52% yield from 3 by treatment with *N*-bromosuccinimide in acetonitrile. The Ullmann homocoupling of 4, facilitated by the presence of aromatic fluorines, gave the desired bis(methoxy) product 5 in x% yield. Demethylation of 5 with BBr₃ furnished F₈BINOL 2 in 88% yield. Finally, recrystallization from methanol/water gave pure 2 as white needles. After several unsuccessful attempts at resolving 2, we were able to chromatographically separate the diastereomeric bis(menthyl)carbonates obtained by reacting racemic 2 with excess (-)-menthylchloroformate. Treatment of each diastereomer with dilute NaOH followed by extraction with diethyl ether afforded (-)-F₈BINOL and (+)-F₈BINOL, respectively. The enantiomeric excess, determined using chiral HPLC (Chiralpak AD column), was found to be >99.9% in each case.

Scheme 1^a

^aKey: (a) *n*-BuLi, ether, -78 °C; (b) 3-methoxythiophene, -78 °C to r.t.; (c) NBS, acetonitrile, r.t.; (d) Cu⁰, 175 °C; (e) BBr₃, dichloromethane, r.t.

Replacement of aromatic hydrogens for fluorines is known to substantially increase barriers to axial torsion in substituted biphenyls. For example, fluorination of the 4 and 5 positions of 9,10-dihydrophenanthrene raises the torsion barrier from 4.1 to 10.3 kcal/mol.⁷ In order to estimate the effect of polyfluorination on atropisomerism in the octafluoro-1,1'-binaphthyl species we prepared racemic 5,6,7,8-octafluoro-1,1'-binaphthyl **6** and determined its X-ray structure.⁸ We chose not to compare the torsion angles in the molecular structures of **1** and **2** due to the possibility of intramolecular OH-F hydrogen bonding in the crystal lattice that could have complicated direct comparison of geometric parameters. Remarkably, the torsion angle between the two tetrafluorinated naphthyl planes in **6** is only 0.7° larger than in the parent hydrido derivative (70.2° for octafluoro-1,1'-binaphthyl vs 69.5° for 1,1'-binaphthyl⁹). To further understand atropisomerism in **2** we investigated acid-promoted racemization of its (-) enantiomer. This process is known to operate for **1**. Remarkably, **2** remains optically active (99.9% e.e) after 24 hours in boiling THF/HCl mixture, whereas **1** rapidly racemizes under these conditions!



6

Polyfluorination of aromatic nuclei is also known to decrease pKa's of bound heteroatoms.⁴ For example, incorporation of four fluorine atoms into the aromatic skeleton of tyrosine results in the pKa' decrease of the ring-bound hydroxyl group by 5 units.¹⁰ We have found that the pKa' of the hydroxyl group in 2 decreases by 1 unit upon octafluorination (1: pKa' 10.28; 2: pKa' 9.29). Another important consequence of fluorination is anodic shift in the oxidation potential of 2, which was found to be more

6

07/21 '99 16:32

667220-2187709

positive than that of binaphthyl by xV, a useful property for applications in oxidation catalysis.

These results lead us to conclude that the effect of fluorine on the reactivity of F₆BINOL will be primarily electronic in nature. The desired conformational flexibility, one of the most important characteristics of BINOL allowing it to coordinate a wide variety of metals, should be preserved. Remarkable configurational stability of either enantiomer of 2 is perhaps its most valuable property. We are currently exploring the scope of the F₆BINOL-derived asymmetric catalysts. The method of the present invention may be used to produce a new Generation of Fluorobinaphthyl Catalysts for several applications including but not limited to: asymmetric catalysis with main group elements, transition metal and lanthanide metals; asymmetric reagent with main group elements, transition metal and lanthanide metals; polymer supported catalysis; nucleophilic displacement of fluorine atoms to modify characteristics of molecule (resulting compounds could be used same manner as original compound); incorporation of molecule into crown ethers for development of phase transfer catalysts; use of compound as a monomer for polymerization; asymmetric polymer supported electrochemical oxidation catalysis; as a chiral auxiliary in an asymmetric reaction; as a resolving agent for chiral compounds, including but not limited to amines; asymmetric catalysis (reagent) in fluorous phase reactions; as a chiral stationary phase for HPLC and other chromatographic techniques; phase transfer catalyst between organic, fluorous phase and alkali solutions.

The foregoing description of the preferred embodiments of the invention has been presented to illustrate the principles of the invention and not to limit the invention to the particular embodiment illustrated. It is intended that the scope of the invention be defined

by all of the embodiments encompassed within the following claims and their equivalents.

REFERENCES

- (1) (a) M. Palucki, N. S. Finney, P. J. Pospisil, M. L. Güler, T. Ishida, E. N. Jacobsen *J. Am. Chem. Soc.* 1998, 120, 948; (b) T. V. Rajanbabu, A. L. Casalnuovo *J. Am. Chem. Soc.* 1996, 118, 6325; (c) A. Schnyder, L. Hintermann, A. Togni *Angew. Chem. Int. Ed. Engl.* 1995, 34, 931;
- (2) R. Noyori *Asymmetric Catalysis in Organic Synthesis*, Wiley: New York, 1994;
- (3) (a) A. S. C. Chan, F. Y. Zhang, C. W. Yip *J. Am. Chem. Soc.* 1997, 119, 4080; (b) F. Y. Zhang, A. S. C. Chan *Tetrahedron: Asymmetry* 1998, 9, 1179; (c) T. Iida, N. Yamamoto, S. Matsunaga, H. G. Woo, M. Shibasaki *Angew. Chem. Int. Ed. Engl.* 1998, 37, 2223; (d) M. Terada, Y. Motoyama, K. Mikami *Tetrahedron Lett.* 1994, 35, 6693;
- (4) B. E. Smart, in: *Organofluorine compounds: Principles and Commercial Applications* (R. E. Banks, ed.), Chapter 3, Plenum Press: New York, 1994;
- (5) methoxythiophene prep;
- (6) BINOL prep via FeCl₃ coupling;
- (7) M. Schlosser, D. Michel *Tetrahedron* 1996, 52, 99 and references cited therein;
- (8) 6 was prepared from 5,6,7,8-tetrafluoro-1-naphthol (G. W. Gribble, C. G. LeHoullier, M. P. Sibi, R. W. Allen *J. Org. Chem.* 1985, 50, 1611) by Ni(0)-catalyzed homocoupling of its trifluoromethanesulfonate ester in NMP at 100 °C;
- (9) R. Kuroda, S. F. Martin *J. Chem. Soc. Perkin Trans II* 1981, 167;
- (10) K. Kim, P. A. Cole *J. Am. Chem. Soc.* 1998, 120, 6851.

THEREFORE WHAT IS CLAIMED IS:

1. A method of producing asymmetric polyfluorinated catalysts, comprising:
fluorinating aromatic groups on binaphthol-based catalysts.
2. A method of producing a family of polyfluoroaryl ligand catalysts,
comprising:
substituting at least one of the hydrogen atoms by fluorine (F) at any one of
the 5,5',6,6',7,7',8,8' positions of 2,2'-dihydroxy-1,1'-binaphthyl (BINOL).
3. A family of polyfluoroaryl ligands, comprising:
2,2'-dihydroxy-1,1'-binaphthyl (BINOL) with at least one of the hydrogen
atoms substituted by fluorine (F) at any one of the 5,5',6,6',7,7',8,8' positions of the
BINOL.
4. The use of the family of polyfluoroaryl ligands comprising 2,2'-dihydroxy-1,1'-
binaphthyl (BINOL) with at least one of the hydrogen atoms substituted by
fluorine (F) at any one of the 5,5',6,6',7,7',8,8' positions of the BINOL as
fluorobinaphthyl catalysts for an application selected from the group
consisting of asymmetric catalysis with main group elements, transition metal and
lanthanide metals, asymmetric reagent with main group elements, transition metal
and lanthanide metals, polymer supported catalysis, nucleophilic displacement of


fluorine atoms to modify characteristics of molecule, incorporation of molecule into crown ethers for development of phase transfer catalysts, use of compound as a monomer for polymerization, asymmetric polymer supported electrochemical oxidation catalysis, as a chiral auxiliary in an asymmetric reaction, as a resolving agent for chiral compounds, including but not limited to amines, asymmetric catalysis (reagent) in fluorous phase reactions, as a chiral stationary phase for HPLC and other chromatographic techniques, and phase transfer catalyst between organic, fluorous phase and alkali solutions.

5. A molecule having the formula,

5,5',6,6',7,7',8,8'-octafluoro-2,2'-dihydroxy-1,1'-binaphthyl (F₈BINOL).

6. The molecule according to claim 5 produced according to scheme I

Scheme I



*Key: (a) *n*-BuLi, ether, -78 °C; (b) 3-methoxythiophene, -78 °C to r.t.; (c) NBS, acetonitrile, r.t.; (d) Cu⁰, 175 °C; (e) BBr₃, dichloromethane, r.t.

//

07/21 '99 16:32

ABSTRACT

The present invention relates to the design and method of synthesis of asymmetric catalysts and more particularly the present invention provides a new generation of polyfluorinated catalysts. Fluorination of aromatic groups on a series of ligands drastically changes their properties including configurational stability and catalytic activity. The fundamental issue is the nature of steric and electronic effects of aromatic fluorination on binaphthol-based catalysts. The basic premise is that alteration of stabilizing stacking and edge-face interactions significantly affects approach of certain substrates to catalytic reaction centers. Due to the high electronegativity of fluorine, electron density in fluoronaphthyl rings is located at the periphery, rather than in the center.